

# **Agriculture and Forestry Climate change report card technical paper**

## **2. Climate and soil functions: impacts on Soil processes and properties and future implications in the UK – a review**

**Mohamed Abdalla & Pete Smith**

Institute of Biological and Environmental Sciences  
University of Aberdeen  
23 St Machar Drive  
Aberdeen AB24 3UU

## Summary

### Current climate

- Soil erosion is an obvious problem in the UK, especially in England and Wales (High confidence).
- Nitrate leaching to the groundwater is high in some localities (High confidence).
- Ammonia emission in the UK is decreasing (Medium confidence).
- Land management reduce greenhouse gas emissions and sequester SOC (GHG) (Low confidence).

### What could happen under climate change?

- Climate change will substantially impact many soil processes and consequently, influence a number of soil properties such as structure (Low confidence), trafficability/workability (Medium confidence), pH (Medium confidence), fertility (Medium confidence) and biota (High confidence).
- Climate change will increase soil erosion (High confidence) and nitrate leaching (High confidence).
- The influence of climate change on soils depends on the size of future changes in climate and on the interaction between the influences of different parameters (Medium confidence).
- Climate change will generally increase GHG emissions (Low confidence) from soils but net primary productivity (NPP) is also expected to increase (Medium confidence).
- Future SOC stock will be determined by the balance between carbon (C) losses from decomposition and C gains from higher plant productivity (Medium confidence).
- SOC stock will significantly be affected by climate-induced changes in land use and management (Medium confidence).
- To increase C sequestration and NPP, and to decrease C flux under climate change, suitable land use and management practices that are resilient to new climatic regimes should be adopted (Medium confidence).

## Introduction

Soils are the foundations on which all terrestrial ecosystems function. They influence water, vegetation and biogeochemical cycling (Rounsevell et al., 1999). Soil chemical and biological processes are controlled by a complex set of factors (Jenny, 1941) but most importantly by the balance between soil temperature and soil moisture. Temperature is key factor that can control many terrestrial biogeochemical processes, including litter (Hobbie, 1996) and soil organic matter (SOM) decomposition, N mineralization and nitrification (MacDonald et al., 1995), denitrification (Abdalla et al., 2009), respiration (Kirschbaum et al., 1995; Christensen et al., 1997; Davidson and Janssens, 2006), CH<sub>4</sub> emissions (Johnson et al., 1996) and plant nutrient uptake (BassiriRad, 2000).

Climate change, due to anthropogenic greenhouse gas (GHG) emissions (IPCC, 2013), is expected to influence soil functions and properties. For the UK, future climate change projections suggest that the mean temperature will increase by 2.4 to 4°C by the year 2080, the maximum and minimum temperature for the winter and summer will increase, and extreme weather events will become more frequent and more severe. Further, summers are expected to be drier and the winter will be wetter, and sea level will rise (UKCP09; Murphy et al., 2009). These changes in temperature and precipitation, in addition to higher atmospheric CO<sub>2</sub> concentrations under climate change, will have substantial impacts on soil functions. Nutrient release from SOM by mineralisation is expected to increase due to higher temperatures (Fang et al., 2005).

Nevertheless, as atmospheric CO<sub>2</sub> concentration and air temperatures increase, soils could lose C to the atmosphere in the form of GHG emissions, resulting in positive feedback that could increase temperature further (Brevik, 2013). Likewise, rainfall affects a number of soil forming processes such as organic matter (OM) turnover, leaching (Stuart et al., 2007) and erosion (McHugh, 2007). Increased winter rainfall due to climate change will increase annual atmospheric N deposition which is currently range from 20 to 30 kg N ha<sup>-1</sup> (Goulding, 1990), if nitrogen emission rates remain constant, but nutrient leaching will also be increased (Liu et al., 2008).

The effect of climate change on soils could occur directly, such as through the effects of temperature on SOM decomposition (Fang et al., 2005) or indirectly, such as through changes in soil moisture, due to changes in plant associated evapotranspiration (Naden and Watts, 2001). Soil moisture is influenced by direct climatic factors (precipitation and temperature), climate induced changes in vegetation, different plant growth rates and changed biogeochemical cycles, different rates of soil water extraction and the effect of enhanced CO<sub>2</sub> levels on plant transpiration (IPCC, 2013).

Soils, especially organic soils, are the largest carbon (C) store in terrestrial ecosystems in the UK (Thomson, 2008). Some important soil functions, like SOM decomposition and nutrient dynamics, are well coupled with plant roots and their related rhizosphere processes (Van Veen et al., 1991). The loss of SOM due to climate change would affect the stability of soil structure, topsoil water holding capacity, nutrient availability, soil erosion and land use, especially in the south of England. Drought will make mineral soils drier, and soils close to low-lying coastal areas could become inundated.

However, soils could have significant role in climate mitigation by sequestering C and reducing the concentration of CO<sub>2</sub> in the atmosphere (Smith, 2012). The sequestration of C in soils could be enhanced by increasing net primary productivity (NPP) relative to C mineralisation or by decreasing C loss from the ecosystem (Six et al., 2004). Improving soil management is important for reducing GHG emissions and for sequestering carbon into SOM (Lal, 2008).

A review of management techniques such as conservation tillage systems (e.g. no-till) in Europe by Holland (2004) found that these practices usually reduce CO<sub>2</sub> fluxes and increase carbon sequestration in the soil compared to conventional tillage systems (Six et al., 2004). West & Post (2002) analysed global data from long-term studies following conversion from conventional to no-tillage. They reported that no-till increased the amount of C sequestration by  $0.57 \pm 0.14 \text{ t C ha}^{-1} \text{ y}^{-1}$  compared with conventional tillage.

However, a recent global review by Abdalla et al. (2013) found significant reductions in CO<sub>2</sub> flux but higher N<sub>2</sub>O emissions compared to conventional tillage. Similarly, Powlson et al. (2014) reported that no-till is beneficial for soil quality and adaptation of agriculture to climate change, but its role in GHG mitigation may have been overstated due to possible change in N<sub>2</sub>O emissions.

The impact of climate change on SOC is still a controversial topic. Two previously published reviews by CLIMSOIL (Schils et al., 2008) and Pacific Northwest National Laboratory (Qafoku, 2014) have found no concrete evidence of climate change effects on SOC. However, to adapt to changes in soil processes, an understanding of how climate and soils interact, and how changes in climate will lead to corresponding changes in soil, is required.

The aim of this review is to investigate the impacts of current and future climate on soil functions of managed agriculture and forestry in the UK. Section 2 focuses on the impacts of climate on soil processes. Section 3 reports on the impacts on soil properties. Section 4

briefly considers the impacts on soil key functions and Section 5 draws some conclusions, and highlights research needs concerning the impacts of climate change on soils in the UK.

### **Impacts on soils processes**

#### *Impacts on Nitrogen (N) mineralisation*

Soil N mineralisation, by which organic matter is converted to inorganic N forms, depends on the amount and nature of the organic matter but also, among others, on temperature and rainfall (Rustad et al., 2001; Rowe et al., 2012). A study of 39 different soil types incubated under standard conditions, showed variable N mineralisation, although the proportion of N mineralised per week was similar (Stanford and Smith, 1972). In mineral and managed agricultural soils, mineralisation rate is correlated directly to soil organic matter content and temperature (Leirós et al., 1999; Abdalla et al., 2009) and soil moisture and rainfall (Emmett et al., 2004). The rate of mineralisation is usually small over the winter when temperature is < 4°C. However, in organic soils, mineralisation results in a large amount of nitrate in the late spring and summer (Defra, 2011).

Changes in rainfall affect soil moisture and thereby, carbon inputs and nitrate uptake by vegetation (Ineson et al., 1998a; Ineson et al., 1998b). In a study covering 665 locations across Britain, Rowe et al. (2012) investigated N availability in relation to variations in climate and reactive N deposition, and found that N mineralisation increased with increasing mean annual temperature of the sites. However, soil characteristics affected this relationship and soil carbon content in particular was a major control on mineralisation rate. They observed that the stock of readily mineralisable N increased more with N deposition in organic compared to mineral soils. This suggested that increasing temperatures due to climate change are likely to increase the effects of N pollution on organic soils more than on mineral soils.

Increasing summer droughts, due to climate change, will reduce N mineralisation (Borken and Matzner, 2009) due to limited soil microorganism activities. However, in saturated soils, lack of oxygen limits N mineralization because only soil microorganisms that can survive under anaerobic conditions are active (Deenik, 2006). Organic rich histosols in agricultural production have higher mineralisation rates for added nitrogenous fertilisers than mineral soils and consequently, higher N<sub>2</sub>O emissions (Skiba et al., 2012).

In a shrubland podzol soil in the UK, net N immobilisation was reported to occur frequently, most likely due to a combination of wetter and colder conditions and low availability of other nutrients (Emmett et al., 2004). Matzner and Borken (2008) reviewed the mechanisms causing the post-frost pulse and suggested that nitrate losses are more likely caused by reduced root uptake rather than by increased N net mineralisation. Net N mineralisation is a complex process and is the difference between gross mineralisation and immobilisation.

Nevertheless, considering all processes increasing or decreasing N production from mineralisation, Ducharme et al. (2007) expected a net increase of 20% in nitrate leaching and nitrate in streams, in France, if appropriate farming practices are not applied. A simulation study by Abdalla et al. (2010) using future climate scenarios in Ireland, predicted that climate change would increase N mineralisation in mineral soils.

#### *Impacts on nitrification and denitrification*

Soil nitrification and denitrification processes play an important role in regulating inorganic N concentration in soils, nitrate leaching and N<sub>2</sub>O emissions. Nitrification is an aerobic process that is directly related to temperature and soil moisture (Emmett et al., 2004) whilst denitrification is an anaerobic process related to soil temperature, water-filled pore space and soil mineral N content (Conen et al., 2000; Sgouridis and Ullah, 2014). Soil ammonia-oxidizing archaea (AOA) are highly abundant and play an important role in the nitrogen cycle (Zhalnina et al., 2012). In acidic forest soil, where net nitrification was high, nitrification was

driven by archaea however, addition of ammonium did not influence the nitrification rate (Stopnisek et al., 2010). In contrast, in nitrogen-rich grassland soils, nitrification was driven by bacteria, not archaea, and the rate of nitrification was related to the abundance of ammonia-oxidizing bacteria (AOB) rather than AOA (Di et al., 2009). Application of ammonium to agricultural alkaline soil increased the abundance of AOB during nitrification process (Jia and Conrad, 2009).

Soil moisture stimulates denitrification by reducing oxygen (Dobbie and Smith, 2001) and increasing solubility of organic carbon and nitrate in soils (Bowden and Bormann, 1986). Future higher atmospheric CO<sub>2</sub> concentration and higher temperature could have strong effects on soil moisture and soil biological activity (Rustad et al., 2001; Zak et al., 2000). Nitrous oxide emissions from soils are mainly regulated by mineral nitrogen, temperature, water content and labile organic compounds (Machefert et al., 2002). Higher mineralisation and denitrification due to increased temperature under climate change will result in higher N<sub>2</sub>O emissions from soils (Abdalla et al., 2010).

The positive relationship between temperature and N<sub>2</sub>O fluxes has been documented in the literature (Flessa and Ruser 2002; Dueri et al. 2007; Abdalla et al., 2014). The higher quantity and frequency of rainfall in late autumn and winter expected in the future will increase denitrification and thereby, N<sub>2</sub>O fluxes from soils (Dobbie et al., 1999). Here, soil moisture increases the supply of C substrate for denitrification (De Catanzaro and Beauchamp, 1985). Nitrous oxide is also produced during nitrate ammonification in soils, but the contribution of this process to the emissions is not well quantified. In this process, NO<sub>3</sub><sup>-</sup> is converted to NO<sub>2</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> (Mohan et al., 2004) and N<sub>2</sub>O is produced when NO<sub>2</sub><sup>-</sup> is reduced (Kelso et al., 1997).

### ***Impacts on decomposition***

Decomposition of litter plays an important role in C cycling in terrestrial ecosystems (Shiels, 2006). Soil respiration, i.e. the total CO<sub>2</sub> efflux at the soil surface, relates to litter decomposition and SOM and comprises autotrophic root respiration and heterotrophic respiration (Bernhardt et al., 2006). Prescott (2009) reported that to sequester more C in soil, more litter should be diverted into humus through microbial and chemical reactions rather than allowing it to decompose. The optimal strategy is to have litter transformed into humic substances and then chemically or physically protected in mineral soil. The addition of N by fertilizer or N-fixing plants is a reasonable way of stimulating humification. Climate change will alter the soil-plant system and impact on decomposition rates, and thereby the amount of SOM. Increasing temperature (Christensen et al., 1997) and atmospheric CO<sub>2</sub> concentration (Ball and Drake, 1998) will increase soil respiration. Elevated CO<sub>2</sub> concentrations will increase photosynthesis, plant growth, belowground C input and substrate and microbial activities (Zak et al., 2000; Anderson et al., 2001) if the soil N is not limited (Hungate et al., 2014).

Increased soil moisture content, due to reduced stomatal conductance and transpiration of plants under high atmospheric CO<sub>2</sub> concentrations, will enhance root and microbial activities and soil respiration (Morgan et al., 2004). However, a global review by Zhang et al (2008) found that a single factor such as climate, litter quality and geographic variable could not, by themselves, explain litter decomposition rates. The combination of litter quality and climatic factors, however, has a very important influence on litter decomposition rates.

Rhizosphere processes are important for the functioning of terrestrial ecosystems. They contribute about 50% of the global CO<sub>2</sub> loss from terrestrial ecosystems (Schimel, 1995), regulate almost all aspects of nutrient cycling (Smith and Read, 1997), and represent the primary gateway for plant water uptake (Jackson et al., 2000). Cheng (2009) found that the magnitude of the rhizosphere priming effect on SOM decomposition varied from 0 to 380% of the unplanted control, and was greatly influenced by plant species and phenology.



However, he suggested a possible decoupling of C cycling with N cycling in the rhizosphere because, the rhizosphere enhancement of soil carbon mineralization did not cause a proportional increase in net N mineralization.

### ***Impacts on ammonia volatilisation***

Agricultural soils account for around 82% of the total ammonia (NH<sub>3</sub>) emissions in the UK. Ammonia emissions predominately arise from livestock manure and urine, though inorganic N fertilisers can also produce NH<sub>3</sub> as nitrogen reacts with compounds in the soil and air (Defra, 2013). This causes fertilizer loss, reduces air quality and causes eutrophication of ecosystems, leading to a loss of biodiversity (Fowler et al., 2013). Ammonia volatilisation is enhanced at higher temperatures and soil drying and therefore, future warmer and drier climates would be expected to increase NH<sub>3</sub> emissions from soils (Skj  th and Geels, 2013). Whitehead et al. (2006) reported higher NH<sub>3</sub> emissions from a groundwater-fed river in southeast England, mainly due to higher temperatures and enhanced microbial activity.

Ammonia can then be nitrified or denitrified further downstream, or re-deposited, increasing N<sub>2</sub>O emissions. However, NH<sub>3</sub> emissions in the UK are decreasing. Projections using the 2010 model structure, in the UK, gave estimates of 306 and 216 Kt NH<sub>3</sub> emissions for the years 1990 and 2020, respectively. This inventory reports emission from livestock agriculture and from nitrogen fertilisers applied to agricultural land (Defra, 2012).

### ***Impacts on nitrate leaching***

Nitrate leaching to the groundwater in the UK is currently high in some localities (Rivett et al., 2007) depending on soil pH (i.e. this is the reason for presence of some nitrate vulnerable zones). Nitrate leaching comes from surplus inorganic N in arable soils during autumn / early winter when the soil is saturated with water (Shepherd et al., 2002) and plant demand for N is low, excessive livestock numbers, inappropriate use of manure and exposure of bare soil during the winter drainage period (Defra, 2000). More than 60% of nitrate leaching to rivers in England has been derived from agricultural land (Defra, 2009).

Nitrate leaching from mineral and organic soils would be almost the same if similar cropping systems and best practice are applied (i.e. about 50 kg N ha<sup>-1</sup>; Stopes et al., 2002). The extent of the leaching depends on rainfall, water holding capacity, soil type, cropping, and the amount / timing of fertiliser or manure applications (White et al., 1983). Nitrate leaching to groundwater depends on the partitioning between run-off and infiltration. Changes in soil hydraulic properties (i.e. the ability of a soil to retain or transmit water and its dissolved constituents) due to higher rainfall intensity will lead to a change in partitioning between run off and recharge.

Climate change will modify soil processes that underpin crop growth, and thereby could increase nitrate leaching in many places in the UK over the next decade (Stuart et al., 2007; Stuart et al., 2011). However, this occurs only under N saturated conditions (Schmidt et al., 2004). Olesen et al (2007) modelled nitrate leaching in the UK for the period 2071 to 2100 and found that nitrate leaching flux has a patchy increase, although this is not quantified. Future nitrate leaching could be mitigated by future changes in agricultural practices, such as planting of catch crops and use of improved crop rotations (Thomsen, 2005; Defra, 2009). However, it may also be influenced by economic responses to climate change (Stuart et al., 2011).

### ***Impacts on soil erosion***

Soil erosion is an obvious problem in the UK, especially in England and Wales, and has significantly increased in recent decades (McHugh, 2007). Soil erosion by water is more widespread than by wind. The UK loses about 2.2 million tonnes of topsoil per year and 17% of the arable land shows signs of erosion due to water driven erosion (EA, 2004). The rate of soil erosion by water has been estimated at 0.1-0.3 t ha<sup>-1</sup> y<sup>-1</sup>. Climate change is likely to

affect soil erosion by water, through its effects on rainfall amount and intensity, soil erodibility, vegetative cover and patterns of land use (McHugh, 2007; Mullan et al., 2011). Climatic variations with future higher winter precipitation intensity (Hulme et al., 1993), and extreme weather events, will create ideal conditions for higher soil erosion. Simulation models predicted that a 10% increase in the winter rainfall in the UK could increase soil erosion by 150% in wet years but the long-term average show modest increase over the current condition (Favis-Mortlock and Boardman, 1995). For England and Wales, McHugh (2007) expected rainfall to increase soil erosion by an average of  $0.6 \text{ t ha}^{-1} \text{ y}^{-1}$  by the 2080s.

For many areas, climate models predict seasonally more intense drying, coupled with increased quantities and intensity of precipitation at other times; conditions that could lead to a large increase in rates of erosion by water. Another potential effect of climate change on soil erosion relates to temperature and  $\text{CO}_2$ -driven changes in plant biomass, with increasing erosion rates possible owing to faster residue decomposition from increased soil microbial activity, i.e. reducing organic matter content and binding of particles (Nearing et al., 2005), or decreasing erosion rates possible with increasing soil surface canopy and biological ground cover (Rosenzweig and Hillel, 1998).

In addition, a more indirect effect of climate change on soil erosion could occur as a consequence of shifting land use and agricultural practice to accommodate the new climatic regime (Williams et al., 1996), i.e. changes in plant biomass and thereby changes in protection of the soil from erosion. Agricultural land will face flooding from rivers and seas. Therefore, some current areas will become unsuitable for agricultural activities due to salt water invading soils and ground water (Barclay, 2012). Generally, soil erosion and structural damage are significantly related to the clearance of natural vegetation for annual cropping and the use of unsuitable farming practices, heavy trampling of soil by sheep and cattle, poor forestry practices and run-off from urban land. Future modifications to planting and harvesting dates, and the implementation of new crops and land use changes are possible, all of which carry the potential to considerably alter rates and patterns of soil erosion (Nearing et al., 2005). However, the introduction of new crops that suit the warmer climate e.g. maize and sunflower which need longer time to provide adequate crop cover and clearance of forests would increase soil erosion problem.

### ***Impacts on soil water and water uptake***

Soil water is controlled by many factors including infiltration, percolation, drainage and run off, and the amount and distribution of rainfall or irrigation (Rounsevell et al., 1999). A decrease in SOM content due to future climate warming (Leirós et al., 1999) would affect soil hydraulic properties (Bowman et al., 2000). Higher future air temperatures will increase evaporation resulting in higher levels of atmospheric water vapour and a greater variability in the amount and intensity of precipitation (Houghton et al. 1992; Kattenberg et al. 1996).

Higher temperatures and accelerating evaporation through the spring lead to drying soils (Hough and Jones, 1997) and will have a great influence on many soil processes and land use. Burt and Shahgedanova (1998) calculated evaporation from 1815 to 1996 for Oxford and reported increases in potential evaporation (PE) but decreases in actual evaporation (AE) due to warmer temperature and less available water during the summer months. Kay et al. (2013) found some evidence of increasing PE throughout the UK since the 1960s. Temperature trends can be used to infer changes in PE as the two are correlated (Dai, 2011; Sheffield et al., 2012), though it is not clear that the relationship will remain constant in a changing climate.

As the result of soil drying and less water uptake by plants, transpiration will reduce under such conditions. However, higher  $\text{CO}_2$  leads to lower stomatal conductance and higher leaf photosynthesis rate (Morgan et al., 2004). Thus, plants in the future will use water more

efficiently to fix atmospheric C than at present. Nevertheless, water shortage, temperature, humidity, vapours pressure and soil nutrients could limit C fixation.

#### *Impacts on seasonal processes*

Climate change will have significant impact on seasonal processes e.g. gross primary productivity (GPP) and respiration, causing disturbances to the terrestrial-atmosphere C-flux balance (Falge et al., 2002). GPP is the main driver of land carbon sequestration. It plays a key role in the global carbon balance and partly offset anthropogenic CO<sub>2</sub> emissions (Janssens et al., 2003). Higher temperatures due to climate change have positive impacts on plant productivity by enhancing the photosynthesis when temperature is in a range of optimum level.

As temperature exceeds the optimum level, it will increase the rate of respiration causing the net ecosystem productivity (NEP) to decline continuously because the increase in respiration is sharper than the increases in GPP (Grace and Zhang (200). A shift in seasonality will result in changing soil C-decomposition, gas fluxes and C sequestration. The dissolved organic carbon (DOC) output from soils results from decomposition of SOM during the dry summer months, which is removed following periods of high rainfall (Dawson et al., 2002). Nevertheless, Freeman et al. (2004) found that increased drying of soils leads to increases in CO<sub>2</sub> production and emissions rather than DOC.

#### **Impacts on soil properties**

##### ***Impacts on soil structure***

Soil structure (aggregate stability and porosity) is considered to be an important soil property and a useful soil health indicator, due to its influential effects on water and gas movement in the soil (Allen et al., 2011). Aggregate stability, the resistance of soil aggregates to external energy such as high intensity rainfall and cultivation, determined by soil structure and a variety of chemical and biological properties and management practices (Moebius et al., 2007; Allen et al., 2011). Soil porosity, a measure of the void spaces in a material as a fraction, controls a range of soil physical indices including soil aeration capacity, plant available water capacity and relative field capacity (Reynolds et al., 2009). Recent studies to model soil water balance and ecosystem conditions under present-day and projected climatic scenarios use porosity as a model parameter (Porporato et al., 2005) because root development and soil enzyme activities are closely related to soil porosity and pore size distribution (Piglai and De Nobili, 1993).

Generally, soil structure is governed by inorganic and organic soil matters, tillage operations and some physical processes (water infiltration, bulk density, rooting depth, and soil surface cover) due to wetting / drying and freeze-thaw conditions (Rounsevell et al., 1999). Soil porosity and aeration status, beside other factors, govern CH<sub>4</sub> (Dalal et al., 2008) and N<sub>2</sub>O (Dalal et al., 2003) gas emissions from soils. Soil structure can also be used to assess soil erosion (Rimal and Lal, 2009). It has a major influence on soil physical properties (Ball, 2013) which indirectly affects GHG production (Gregorich et al., 2006). Structure can override the influence of texture in regulating gas exchange, mainly because of its substantial influence on soil water content and pore continuity in soils of the same type. Soil structure affects roots ability to grow and supply leaves with water and nutrients (Passioura, 1991).

In adverse situation, it induces them to send hormonal signals that slow the growth of the shoot, even if they are able to take up sufficient water and nutrients (Passioura, 1991). These effects of soil structure on plant growth will also affect GHG emissions from soils. Climate change could influence soil structure by modifying soil physical processes and the amount of SOC available in soils (Carter and Stewart, 1996). Future low summer precipitation will lead to soil shrinkage and increased cracking, especially in clay-rich soils (Harrison et al., 2012). In a study in north Wales, Dominguez et al (2015) found that soil



respiration in podzolic (organo-mineral) soils from wet shrublands was more vulnerable to recurrent drought than to warming, and that the drought impact did not diminish at decadal time scales. This stimulation of soil respiration was due to changes in soil structure which led to 54% reduction in water holding capacity. They suggested that changes in sub-dominant vegetation and soil physical properties may determine the climate change impacts on soil C dynamics.

Geris et al. (2015) examined short-term effects of extreme drought on storage dynamics and runoff response in hydopedological units in a headwater catchment in the Scottish highlands using isotopes experiments. They found incredible small storage changes in histosols compared with those in moorland and forested podzols. They suggested that during dry periods, large parts of the catchment are disconnected from the river network and run-off is generated mainly from the wet histosols however, during events, there was an intermittent connection of the hillslopes that recharged the wetland and stream. This participated to recovery and resilience of the catchment in its run-off response.

#### *Impacts on soil pH*

Soil pH is the degree of soil alkalinity or acidity and has a great impact on nutrient availability and micro-organism activity in soil. Changes in rainfall amount and intensity, and increased temperature due to climate change, will influence leaching intensity and soil mineral weathering, and thereby soil pH. In a European simulation study on forestry soils, Reinds et al. (2009) reported that climate change would lead to higher weathering rates and nitrogen uptake and limited positive effects on recovery from acidification compared to current climate.

Evans et al. (2005) used the MAGIC dynamic model (Model of Acidification of Groundwater In Catchments) and climate scenarios of rising temperature, and decreasing rainfall in the UK, to estimate impacts of climate change on soil and water recovery from acidification. They reported that high dissolved organic carbon (DOC) and elevated organic acidity, due to higher temperature, is expected to lower soil pH and increase leaching of basic cations into surface waters, bringing about recovery of these waters from acidity. However for agricultural soils, under climate change, lime could be used to control acidity. Soils with high clay and organic matter content are more able to resist a drop or rise in pH whilst sandy soils are more vulnerable to acidification.

Clay content cannot be modified, but organic matter content can be changed by following best land management practices. Salinization due to weather extremes e.g. drought and rainfall periods under climate change could inhibit biological N transformation (Curtin et al., 1999), N fixation capacity by legumes (Delgado et al., 1993) and decrease plant growth (i.e. less N use efficiency and higher N loss in the form of gases and leaching). Further, a combination of high pH and sodium has detrimental impacts on soil properties with implications for ecosystem function and services. This combination is associated with colloid dispersion, loss of organic carbon, decrease in soil permeability, and increase in run-off and erosion (Defra, 2011). However, salinity due to higher evaporation rates in relation to irrigation applied is not considered a likely problem in the UK, even under climate change (Defra, 2010).

#### *Impacts on traffic-ability/ workability*

Trafficability is the capability of soil to permit movement of a vehicle over the land surface whilst workability the number of days in a given period suitable for field work (Reeve and Fausey, 1974). Climate change will affect both field traffic-ability and workability (Campbell and O'Sullivan, 1991). Soil moisture status is the most important influencing factor in determining the traffic-ability / workability of a land. Wet soils, during critical periods for management operations such as harvesting and ploughing will limit machinery access, and soils could be subject to compaction and structural damage (Earl, 1997). Reduction in

rainfall could improve soil workability in some wet and heavy soil areas in Scotland (MacDonald et al., 1994). Projected climate change scenarios showed a small increase in the number of workable days in the UK, due to lower soil water content at higher temperatures. However, localised threat of soil compaction will remain (Defra, 2010). A recent simulation study by Harding et al. (2015) suggests that future field operations across the UK will start earlier due to the dramatic decrease in frosts, especially in the highlands. Thus, growing seasons will be longer, which may increase crop productivity and carbon returns to the soil.

### ***Impacts on soil biota***

Soil is a haven for a huge diversity of bacteria, fungal species and animals. These organisms make up soil food web which is essential for the functioning of terrestrial ecosystems as its key role is to recycle unused organic matter derived from the above-ground (Bardgett 2005). Changes in future rainfall, including drought and flooding, will directly impact soil biota through changing soil water, and indirectly through changing the soil habitat e.g. shrinkage and swelling of clay rich soils (Harrison et al., 2012). Drought could result in less microbial biomass, less microbial activity, and the death of some larger soil organisms (Gordon et al., 2008; De Vries and Shade, 2013).

Climate change can also impact soil biota by increasing soil erosion, especially where extreme events increase and where climate-change induced changes in land use and management increase soil vulnerability to erosion (Nearing et al., 2004). A recent review by Blankinship et al. (2011) investigated the impacts of climate change on soil biota and reported that microbial biomass is significantly increased by elevated CO<sub>2</sub>, but bacterial abundance is negatively affected by warming, and fungal biomass increased with increasing precipitation. Elevated CO<sub>2</sub> leads to higher plant photosynthesis and growth, and thereby increasing the rates of carbon input to soil, which in turn strongly modifies the growth and activity of soil biota (Phillips et al., 2011; Phillips et al., 2012).

Warming of the climate could stimulate the exudation of carbon from plant roots (Yin et al., 2013) and thus microbial feedback to climate change (Grayston et al., 1996); it increasing bacterial and fungal biomass, and affects the structure of food webs (De Vries and Bardgett, 2014). Nielsen et al. (2011) reported that changes number of functional groups present in soil and the diversity of these groups can result in significant effects on ecosystem functioning. Further, the effects of climate change on the composition and functioning of soil food webs could result in changing plant communities under the new condition (De Vries et al., 2014). However, changes in plant communities might be more important than direct effects of climate change for ecosystem functioning (Ward et al., 2013).

### ***Impacts on soil fertility***

The availability of soil nutrients has a great influence on plant growth and water use. Soil fertility is essentially related to SOM (Brevik, 2013). Climate change could cause soil degradation through erosion and losses of SOM. Low SOM content decreases soil fertility as a result of low nutrient and reduced water holding capacity, leading to a deterioration of soil structure (King et al., 2005). Higher mineralisation under climate change will also increase N loss by ammonia volatilisation (Rounsevell et al., 1996). Mineral and drained organic soils with sufficient amounts of OM are more productive than soils that have low organic matter.

Agriculture contributed 26% of the English total phosphorus (TP) load, 22% of the Scottish TP load, 57% of the Welsh TP load and about 47% of the TP load to inland and coastal waters of Northern Ireland (Defra, 2006). Agricultural soils are rich with P because of application of phosphate fertilizer for many years (Donnison, 2011). Intensively managed grassland could represent a source for reactive phosphate due to manure application and accumulation of P. The principle pathway for P to enter water is by erosion and overland flow. Enrichment of an ecosystem with N and P leads eutrophication that causes algae and

higher forms of plant life in water to grow too fast and stimulate the growth of certain on land plants. This can disturb the balance of organisms present in the water and the quality of the water concerned and make some on land plants dominant so that the natural diversity would be lost (Defra, 2011).

In a climate change simulation study for north England, Bouraoui et al. (2002) found higher nutrient uptake and high crop growth; however, nutrient (N and P) losses to surface water were increased due to climate accelerating soil processes such as mineralisation of SOM, and increasing water loss from the soil profile to rivers in case of extreme rainfall events. They expected great impacts on soils and crop management, and suggested a need for adjusting current management practices. To decrease N and P losses management of permanent pasture should be improved by reducing stocking density, nutrient input, run-off and erosion (Defra, 2009).

### **Impacts on the soil key functions**

#### *Impacts on SOC stock*

The relationship between soil functions and SOC is strong (Tate, 1992). Changes in SOC take place over a long period of time and are governed by many factors such as climatic (e.g. temperature and precipitation) and edaphic (e.g. soil parent material, clay content, cation exchange capacity) factors (Dawson et al., 2007). Chapman et al. (2013) reported no change in the overall total soil C stock at 100 cm depth, for 25 years period, across the Scotland. However, they found C stock for soils under woodland (mainly coniferous plantation) significantly increased. By recalculating the C stock to a depth of 15 cm the overall C stock (when deep peat sites were excluded) was significantly increased. Though both improved grassland soils and those initially under arable cultivation showed a significant decrease in C content. Another study by Reynolds et al. (2013) covering a 30 years' period (1978-2007) also reported no change in soil C across Great Britain for the depth of 15cm. Further, Hopkins et al. (2009) found no consistently significant changes in SOC stocks due to climate over a long-term (>100 years) experiment on grassland soil.

Bellamy et al. (2005) reported that climate change led to SOC loss from England and Wales at a rate of 0.6% per year over the period between 1978 and 2003. However, Smith et al. (2007) found that the main driver for C loss is land use change and climate change was responsible for only 10-20% of the amount of SOC loss reported by Bellamy et al. (2005). Using isotope experiments, Evans et al. (2007) also reported these negative impacts of agriculture intensification in the UK on the loss of SOC. By reanalysing the data of England and Wales (1978-2003), Kirk and Bellamy (2010) suggested that modifications in land use (conversion of natural vegetation and grassland to arable land) and improved management practices (drainage, mineral fertilizers and stocking rates) were the most significant drivers for the C loss than climate change.

More recently, the analysis of Barraclough et al. (2015) suggested that only 0-5% of soil C change in mineral soil could be attributed to climate change, though 9-22% of soil C change in organic soils could be due to climate change. However, although there is a conflict between the evidences regarding the impacts of climate change on SOC in forests and grasslands, consistent evidences show reduction of SOC in arable agriculture due to climate change (Goidts and Van Wesemael, 2007; Chapman et al., 2013). Ciais et al. (2010) reported that management and land use will have more significant impacts on future SOC than the climatic factors. Rees et al. (2005) found that the increases of crop yield under climate change did not guarantee an increase in SOC. They observed an increase of 4.26 t ha<sup>-1</sup> in wheat yield between 1948 and 2001 but this did not prevent SOM loss (Jastrow et al., 2007). With climate change, the NPP and litter input will increase and could compensate for the SOC loss (Smith et al., 2006; Gottschalk et al., 2012). Thus, the amount of SOC depends on the balance between the impacts of future increased temperature and

decreased soil moisture on decomposition rates and C losses from decomposition and C gains from higher plant productivity (Smith, 2012).

Fluvial C fluxes also represent an important part of the C budget; they constitute an important pathway for carbon loss from organic soils. Soil C losses increase through losses of DOC and particulate organic carbon (POC). DOC is the carbon included within SOM in solution. Schulze and Freibauer (2005) reported that DOC loss had originated from the SOM, therefore contributing to the overall decrease in SOC. The loss of DOC is controlled by rainfall but also, among others, is associated with solar radiation and temperature (Harrison et al., 2008).

Long-term increase in DOC has a severe negative impact on soil biota, water industry (i.e. increase the cost) and C stock (Evans et al., 2005). McCartney et al. (2003) found a clear rising trend in DOC concentrations in a stream draining a mature forest catchment in Scotland from around 5 mg l<sup>-1</sup> in the early 1980s to around 16 mg l<sup>-1</sup> in 2003. Climate change is expected to influence the loss of DOC from soils to rivers and lakes in the UK. Freeman et al. (2001) reported an increase of 65% in DOC concentration in freshwaters draining from 20 sites, of different soils, land uses and locations in the UK, and concluded that rising temperatures increased DOC loss from soils. Evans et al. (2005) reported an increase in DOC by 91% in the UK water over 15 years period. However, in a long-term study, Worrall et al. (2003) found that the increases in DOC concentrations in rivers coincided with increases in mean summer temperatures. Worrall et al. (2004) suggested that DOC is most likely driven by temperature and the frequency of severe droughts. Worrall and Burt (2004) confirmed that climate, especially severe drought, is a main driver for DOC loss.

Pawson et al (2008) reported that POC accounted for 80% of a very large aquatic C flux of 93g C m<sup>-2</sup> y<sup>-1</sup> in the south Pennine catchment. Oxidation of this POC amount will create a great C loss in the form of CO<sub>2</sub> (Schlesinger, 1995). However, Smith et al. (2001) argue that only a small fraction of the eroded POC is decomposed and released as CO<sub>2</sub> to the atmosphere. Changes in rainfall due to climate change will influence runoff, which drives POC (Dawson et al., 2002). Furthermore, Chaplot and Cooper (2015) noted a tendency for clayey soils, which were fully covered by grass, to present stable aggregates and thus to yield greater CO<sub>2</sub> emissions, but lower POC and DOC outputs, than degraded sandy soils of low aggregate stability.

### ***Impacts on greenhouse gas emissions***

Production and release of GHGs are essentially due to biological processes; however, soil physical condition can impact biology by affecting the soil physical environment (Gregorich et al., 2006). Changes in temperature (Fiscus et al., 1997) and precipitation (Izaurrealde et al., 2003; Mearns, 2003) due to climate change will influence mineralization and denitrification, and thereby GHG production. The loss of SOM under high temperature will reduce soil fertility and consequently deteriorate soil structure. Higher temperatures influence microbial activity in soils and increase GHG emissions to the atmosphere (Ball, 2013). Climate change would reduce porosity and soil aeration resulting in reductions in rates of CH<sub>4</sub> oxidation and CO<sub>2</sub> release from decomposition, but increasing rates of denitrification and N<sub>2</sub>O release. Smith et al (2003) found that gas diffusivity was very important for CH<sub>4</sub> oxidation rate. Well-aerated, moist soil is suitable for CH<sub>4</sub> oxidation and CO<sub>2</sub> release (Ball, 2013).

However, higher temperature alone could reduce emissions of CH<sub>4</sub> by changing the hydrology of the soil. Temperature increases evapotranspiration and thereby lowers the water table leading to decreased CH<sub>4</sub> emissions, since aerobic conditions enhance CO<sub>2</sub> release relative to CH<sub>4</sub> (Moore et al., 1998). Moreover, changes in rainfall could lead to changes in the water table, and thus CH<sub>4</sub> emissions from soils (Mojeremane et al., 2010). Nitrous oxide production and emissions in soils is stimulated by temperature and water filled pore space (WFPS; an indicator for soil aeration status) (Smith et al., 2003). Dobbie and



Smith (2001) found a linear relationship between WFPS and mean N<sub>2</sub>O fluxes for a grassland soil in Scotland.

Land management with climate change (i.e. climate-induced changes in land management) can have substantial impacts on soils. Graves and Morris (2013) found interaction between the projected land use management scenarios and climate change resulted in soil degradation. Schils et al. (2008) estimated the loss of SOC in Europe due to land use change especially drainage, by 20-40 tonnes of CO<sub>2</sub> ha<sup>-1</sup>y<sup>-1</sup>. Lowering water table in organic soils by drainage increases CO<sub>2</sub> loss but decreases CH<sub>4</sub> emissions to the atmosphere (Smith et al., 2003). However, restoration could reduce C fluxes, but may increase CH<sub>4</sub> emissions (e.g. Waddington and Day, 2007). Precision application of N (Pierce and Nowak, 1999) and management to reduce the fallow period and/ or minimise soil disturbances reduce N<sub>2</sub>O losses and thereby, GHG emissions (Osborne et al., 2010). Webb et al. (2001) reported that the loss of mixed farming will result in soil C loss.

This is because the association between crop and animal outputs can give suitable inputs of nutrients to soils and protect the crop. The decrease in SOC content in the English arable soils is due to replacing grassland in mixed farming rotations by permanent arable cropping (King et al., 2005). In a literature review, Post and Kwon (2000) reported that land use change from arable cropping to grassland would increase soil C by 33 g C m<sup>-2</sup> y<sup>-1</sup> however, rainfall intensity and the species sown in the new pasture can considerably affect the rate. Tillage practices and climate variations affect the release of GHGs from soils. No-tillage increases N<sub>2</sub>O emissions, but decreases CO<sub>2</sub> emissions (Ball et al., 1999; Abdalla et al., 2013). Changes in vegetation cover could alter runoff and nutrient losses as well as SOM content through C input to soils. Soil ploughing increases CO<sub>2</sub>-evolution (Reicosky and Archer, 2007) because ploughing increases soil disturbance, crop residues distribution (Grigera et al., 2007) and microclimate i.e. increases soil temperature (Vinther and Dahlmann-Hansen, 2005). However, an application of combined management to reduce N and C emissions could be a useful approach to prevent trade off and swapping of emissions between the GHG gases CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O (Schils et al., 2008).

Holman et al. (2005) predicted a possible future reduction in the arable land area in East Anglia and conversion of upland grass to arable in the Northwest as a result of economic pressure under future climate scenarios. They also predict the introduction of new crops such as sunflower, grain and forage maize and the altering of current rotations. However, soil erosion and loss of SOM have the potential to limit the ability of farmers to take advantage of future opportunities to increase agricultural production (MLCC, 2013). Preserving upland vegetation cover is a key win-win management strategy that will reduce erosion and loss of soil carbon, and protect a variety of services, such as the continued delivery of a high quality water resource.

Further, the growing of energy crops such as willow and *Miscanthus* which increase SOM and mitigate GHG emissions (Hiller et al., 2009) could be encouraged under future climatic condition (Evans et al., 1995). Land use change will also be moderated by potential policy goals that seek to reduce GHG emissions from land and / or increase the size of land-based sinks. This will include strategies to reduce C and N fluxes through increased efficiency, afforestation and biomass production.

## Conclusions

We have reviewed the impacts of anthropogenic climate and atmospheric CO<sub>2</sub> concentration change on soil functions in the UK. Currently, soil erosion is high especially in England and Wales, nitrate leaching is high in some localities but ammonia emission is decreasing. Also, arable agriculture is currently losing SOC. Climate change will accelerate soil processes, and could lead to more rapid decomposition of SOM, increased microbiological activity, quicker release of nutrients, increased rates of nitrification and denitrification, increased



nitrate leaching and soil erosion. These changes in soil processes will influence soil structure, pH, trafficability / workability and fertility. However, the extent of this influence depends on the size of future changes in climate and on the interaction between the influences of different parameters. Climate change would reduce CH<sub>4</sub> oxidation and CO<sub>2</sub> release from decomposition, but increase rates of denitrification and N<sub>2</sub>O release. Under climate change, the NPP and litter input will increase and could compensate for potential SOC losses due to increased decomposition rates.

The net SOC stock will therefore, be determined by the balance between the C losses from decomposition and C gains from the high crop productivity. However, this will significantly be affected by the climate-induced changes in land use and management. Most of the available evidences show that management and land use systems and not climate change will have the greater impacts on future SOC. However, the evidences from experimental and simulation researches on soils and impacts of future climate change on SOC in the UK are inadequate. To fill this key knowledge gap we suggest proposing new strategies that emphasize additional necessary researches.

## References

- Abdalla M, Wattenbach M, Smith P, Ambus P, Jones M, Williams M (2009) Application of the DNDC model to predict emissions of N<sub>2</sub>O from Irish agriculture. *Geoderma* 151:327-37.
- Abdalla, M., Jones, Williams, M (2010) Simulation of nitrous oxide emissions from Irish arable soils: effects of climate change and management. *Biology and Fertility of Soils* 46:247-260.
- Abdalla M, Osborne B, Lanigan G, Forristal D, Williams M, Smith P, Jones MB (2013) Conservation tillage systems: a review of its consequences for greenhouse gas emissions. *Soil Use & Management* 29:199-209.
- Abdalla M, Hastings A, Helmy M, Prescher A, Osbourne B, Lanigan G, Forristal D, Killi D, Maratha P, Williams M, Rueangritsarakul K, Smith P, Nolan P, Jones MB (2014) Assessing the combined use of reduced tillage and cover crops for mitigating greenhouse gas emissions from arable ecosystem. *Geoderma* 223, 9-20.
- Allen DE, Singh BP, Dalal RC (2011) Soil Health Indicators, Soil Health and Climate Change: A Review of Current Knowledge. In B.P. Singh et al. (eds.), *Soil Health and Climate Change*, Soil Biol. 29.Springer-Verlag, Berlin 25-45.
- Anderson LJ, Maherali H, Johnson HB, Wayne H, Jackson RB (2001) Gas exchange and photosynthetic acclimation over subambient to elevated CO<sub>2</sub> in a C3-C4 grassland. *Global Change Biology* 7: 693-707.
- Ball BC (2013) Soil structure and greenhouse gas emissions: a synthesis of 20 years of experimentation. *European Journal of Soil Science* 64: 357-373.
- Ball AS, Drake BG (1998) Stimulation of soil respiration by carbon dioxide enrichment of marsh vegetation. *Soil Biology and Biochemistry* 30: 1203-1205.
- Ball BC, Scott A, Parker JP (1999) Field N<sub>2</sub>O, CO<sub>2</sub> and CH<sub>4</sub> fluxes in relation to tillage, compaction and soil quality in Scotland. *Soil Tillage Research* 53:29-39.
- Barclay C (2012) *Agriculture and Climate Change*. House of the common library. SN/SC/3763. Science and Environment Section.

- Bardgett RD (2005) *The Biology of Soil: A Community and Ecosystem Approach*. Oxford Univ. Press, Oxford.
- Barraclough D, Smith P, Worrall F, Black H, Bhogal A (2015) Climate change and soil carbon declines in England and Wales 1978-2003. *European Journal of Soil Science* 66: 451-462.
- BassiriRad H (2000) Kinetics of nutrient uptake by roots: responses to global change. *New Phytologist* 147: 155-169.
- Bellamy PH, Loveland PJ, Bradley RI, Lark RM, Kirk GJD (2005) Carbon losses from all soils across England and Wales 1978-2003. *Nature* 437: 245-248.
- Bernhardt ES, Barber JJ, Phippen JS, Taneva L, Andrews JA, Schlesinger WH (2006) Long-term effects of Free Air CO<sub>2</sub> Enrichment (FACE) on soil respiration. *Biogeochemistry* 77: 91-116.
- Blankinship J, Niklaus P, Hungate B (2011) A meta-analysis of responses of soil biota to global change. *Oecologia* 165:553-565.
- Borken W, Matzner E (2009) Reappraisal of drying and wetting effects on C and N mineralization and fluxes in soils. *Global Change Biology* 15:808-824.
- Bowman RS, Sullivan EJ, Li Z (2000) Uptake of cations, anions, and nonpolar organic molecules by surfactant-modified clinoptilolite-rich tuff. p. 287-297. In C. Colella and F. A. Mumpton (eds.) *Natural zeolites for the third millennium*. De Frede Editore, Naples, Italy.
- BowdenWB, Bormann FH (1986) Transport and loss of nitrous oxide in soil water after forest clear cutting. *Science* 233: 867-869.
- Bouraoui F, Galbiati L, Bidoglio G (2002) Climate change impact on nutrient loads in the Yorkshire Ouse catchment (UK). *Hydrol. Earth System Science* 6: 197-209.
- Brevik EC (2013) *An Introduction to Soil Science Basics*. In *Soils and Human Health*; Brevik EC, Burgess LC, Eds.; CRC Press: Boca Raton, FL, USA, 2013; pp. 3-28.
- Burt TP, Shahgedanova M (1998) An historical record of evaporation losses since 1815 calculated using long-term observations from the Radcliffe Meteorological Station, Oxford, England. *Journal of Hydrology* 205: 101-111.
- Campbell DJ, O'Sullivan LA (1991) 'The Cone Penetrometer in Relation to Trafficability, Compaction, and Tillage' in Smith, K.A. and Mullins, C.E. *Soil Analysis: Physical methods*, Marcel Dekker, New York.
- Carter MR, Stewart BA (eds) (1996) *Structure and Organic Matter Storage in Agricultural Soils*. CRC Press, Boca Raton, Fla., USA. 477pp.
- Chaplot V, Cooper M (2015) Soil aggregate stability to predict organic carbon outputs from soils *Geoderma* 243-244: 205-213.
- Chapman SJ, Bell JS, Campbell CD et al. (2013) Comparison of soil carbon stocks in Scottish soils between 1978 and 2009. *European Journal of Soil Science* 64: 455-465.
- Cheng W (2009) Rhizosphere priming effect: Its functional relationships with microbial

- turnover, evapotranspiration, and C-N budgets. *Soil Biology and Biochemistry* 41:1795-1801.
- Christensen TR, Michelsen A, Jonasson S, Schmidt IK (1997) Carbon dioxide and methane exchange of a subarctic heath in response to climate change related environmental manipulations. *Oikos* 79:34-44.
- Ciais P, Wattenbach M, Vuichard N, Smith P, Piao SL, Luyssaert S, Janssens I, Bondeau A, Dechow R, Smith PC, Beer C, van der Werf GR, Gervois S, Van Oost K, Tomelleri E, Don A, Freibauer A, Schulze ED, Abril G, Bouriaud O, Churkina G, Grace J, Jung M, Nabuurs GJ, Paris JD, Papale D, Peylin P, Reichstein M, Schellhaas MJ, Soussana JF, Vetter M, Viovy N, Zaehle S (2010) The European Carbon Balance. Part 2: Croplands. *Global Change Biology* 16: 1409-1428.
- Conen F, Dobbie KE, Smith KA (2000) Predicting N<sub>2</sub>O emissions from agricultural land through related soil parameters. *Global Change Biology* 6: 417-426.
- Curtin D, Steppuhn H, Campbell C A, Biederbeck VO (1999) Carbon and nitrogen mineralization in soil treated with chloride and phosphate salts. *Canadian Journal of Soil Science* 79: 427-429.
- Dai A (2011) Drought under global warming: A review. *WIREs Climate Change* 2: 45-65.
- Dalal RC, Eberhard R, Grantham T, Mayer DG (2003) application of sustainability indicators, soil organic matter and electrical conductivity, to resource management in the northern grains region. *Australian Journal of Experimental Agriculture* 43:253-259.
- Dalal R C, Allen D E, Livesley S J and Richards G (2008) Magnitude and biophysical regulators of methane emission and consumption in the Australian agricultural, forest, and submerged landscapes: a review. *Plant and Soil* 309: 43-76.
- Davidson EA and Janssens IA (2006) Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440:165-173.
- Dawson JJC, Billett MF, Neal C, Hill S (2002) A Comparison of Particulate, Dissolved and Gaseous Carbon in Two Contrasting Upland Streams in the UK. *Journal of Hydrology* 257: 226- 246.
- Dawson JJC, Godsiffe EJ, Thompson IP, Ralebitso-Senior TK, Killham KS, Paton GI (2007) Application of biological indicators to assess recovery of hydrocarbon impacted soils. *Soil Biology and Biochemistry* 39:164-177.
- De Catanzaro JB, Beauchamp EG (1985) The effect of some carbon substrates on denitrification rates and carbon utilization in soil. *Biology and Fertility in Soils* 1:183-187.
- Deenik J (2006) Nitrogen Mineralization Potential in Important Agricultural Soils of Hawai'i. *Soil and Crop Management SCM*-15.
- Defra (2006) Estimate of the sources of phosphorus in UK waters. White PJ, Hammond JP (2006). A Defra funded project WT0701CSF.
- Defra (2010) Use of 'UKCIP08 Scenarios' to determine the potential impact of climate change on the pressures/threats to soils in England and Wales. Cooper D, Gooday R, Hallett P, Irvine B, Morrow K, Ragab R, Rawlins B, Richards M, Smith P, Tye A. Defra Project SP0571

- Defra (2011) Evaluation of the Occurrence of Soils with pH Higher than 8.3 Observed Within the Countryside Survey. Lebron I, Keith AM, Hughes S, Robinson DA, Reynolds B, Cooper D, Emmett BA(2011). Project SP1304. Centre for Ecology and Hydrology, UK.
- Defra (2013) Department for Environment, Food and Rural Affairs 2013. Air pollution in the UK.
- Delgado MJ, Garrido JM, Ligeró F, Lluch C ( 1993 ) Nitrogen fixation and carbon metabolism by nodules and bacteroids of pea plants under sodium chloride. *Physiologia Plantarum* 89: 824-829.
- De Vries FT, Bardgett RD (2014) Terrestrial Biodiversity Climate change Impacts Report Card. Technical paper, Climate change effects on soil biota in the UK.  
<http://www.manchester.ac.uk/research>
- De Vries FT, Shade A (2013) Controls on soil microbial community stability under climate change. *Frontiers in Microbiology* 4:265.
- Di HJ, Cameron KC, Shen JP, Winefield CS, O'Callaghan M, Bowatte S, He JZ (2009) Nitrification driven by bacteria and not archaea in nitrogen-rich grassland soils. *Nature Geoscience* 2: 621-624.
- Dobbie KE, Smith KA (2001) The effects of temperature, water filled pore space and land use on N<sub>2</sub>O emissions from an imperfectly drained gleysol. *European Journal of Soil Science* 52: 667-673.
- Dominguez MT, Sowerby A, Smith AR, Robinson DA, Van Baarsel S, Mills RTE, Marshall MR, Koller E, Lebron I, Hall J, Emmett BA (2015) Sustained impact of drought on wet shrublands mediated by soil physical changes. *Biogeochemistry* 122:151-163.
- Donnison L (2011) Review of the effects of farmland trees on erosion and pollution in the local farmed landscape. Report to Woodland Trust.
- Ducharne A, Baubion C, Beaudoin N, Benoit M, Billen G, Brisson N, et al (2007) Long term prospective of the Seine River system: confronting climatic and direct anthropogenic changes. *Science of the Total Environment* 375:292-311.
- Dueri S, Calanca PL, Fuhrer J (2007) Climate change affects farm nitrogen loss-A Swiss case study with a dynamics farm model. *Agricultural Systems* 93:191-214.
- EA (2004). Environment Agency (2004). The state of soils in England and Wales.
- Earl R (1997) Prediction of trafficability and workability from soil moisture deficit. *Soil and Tillage Research* 40:155-168.
- Emmett BA, Beier C, Estiarte M, Tietema A, Kristensen HL, Williams D, Penuelas J, Schmidt I, Sowerby A (2004) The response of soil processes to climate change: results from manipulation studies of shrublands across an environmental gradient. *Ecosystems* 7:625-37.
- Evans LG, Eckersten H, Semenov MA, Porter JR (1995) Effects on willow. In: Harrison, PA, Butterfield RE, Downing TE (Eds.), *Climatic Change and Agriculture in Europe*.

Assessment of Impacts and Adaptations. University of Oxford, UK, pp. 220-223 Research Report No. 9. Environmental Change Unit.

Evans CD (2005) Modeling the effects of climate change on an acidic upland stream, *Biogeochemistry* 74: 21-46.

Evans CD, Monteith DT, Cooper DM (2005) Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts. *Environmental Pollution* 137: 55-71.

Evans CD, Freeman C, Cork LG, Thomas DN, Reynolds B, Billett MF, Garnett MH, Norris D (2007) Evidence against recent climate-induced destabilisation of soil carbon from C-14 analysis of riverine dissolved organic matter. *Geophysical Research Letters* 34: L07407.

Falge E, Baldocchi D, Tenhunen J, Aubinet M, Bakwin P, Berbigier P, Bernhofer C, Burba G, Clement R, Davis KJ (2002) Seasonality of ecosystem respiration and gross primary production as derived from FLUXNET measurements. *Agricultural and Forest Meteorology* 113:53-74.

Fang C, Smith P, Moncrieff J B, Smith J U (2005) Similar response of labile and resistant soil organic matter pools to changes in temperature. *Nature* 433: 57-59.

Favis-Mortlock DT, Boardman J (1995) Nonlinear responses of soil erosion to climate change: a modelling study on the UK South Downs. *Catena* 25: 365-387.

Flessa H, Ruser R (2002) N<sub>2</sub>O and CH<sub>4</sub> fluxes in potato fields: automated measurement, management effects and temporal variation. *Geoderma* 105:307-325.

Fowler D, Pyle JA, Raven JA, Sutton MA (2013) The global nitrogen cycle in the twenty-first century: introduction. *Philosophical Transactions of the Royal Society B* 2013 368 20130120; DOI: 10.1098/rstb.2013.0120.

Freeman C, Evans CD, Monteith DT, Reynolds B, Fenner N (2001) Export of organic carbon from peat soils. *Nature* 412: 785.

Freeman C, Fenner N, Ostle NJ, Kang H, Dowrick DJ, Reynolds B, Lock MA, Sleep D, Hughes S and Hudson J (2004) Export of dissolved organic carbon from peatlands under elevated carbon dioxide levels. *Nature* 430:195-198.

Fiscus EL, Reid CD, Miller JE, Heagle AS (1997) Elevated CO<sub>2</sub> reduces O<sub>3</sub> flux and O<sub>3</sub>-induced yield losses in soybeans: possible implications for elevated CO<sub>2</sub> studies. *Journal of Experimental Botany* 48:307-13.

Geris J, Tetzlaff D, Soulsby C (2015) Resistance and resilience to droughts: hydrological controls on catchment storage and run-off response. *Hydrological Processes*. Published online in Wiley Online Library (wileyonlinelibrary.com) DOI: 10.1002/hyp.10480.

Goidts, E., van Wesemael, B., 2007. Regional assessment of soil organic carbon changes under agriculture in Southern Belgium (1955-2005). *Geoderma* 141: 341-354.

Gordon H, Haygarth PM, Bardgett RD (2008) Drying and rewetting effects on soil microbial community composition and nutrient leaching. *Soil Biology and Biochemistry* 40:302-311.



- Gottschalk P, Smith JU, Wattenbach M, Bellarby J, Stehfest E, Arnell N, Osborn T, Jones C, Smith P (2012) How will organic carbon stocks in mineral soils evolve under future climate? Global projections using RothC for a range of climate change scenarios. *Biogeosciences* 9: 3151-3171.
- Goulding K W T (1990) Nitrogen deposition to land from the atmosphere. *Soil Use and Management* 6:61-63
- Graves AR, Morris J (2013) Restoration of fenland peatland under climate change. Report to the Adaptation Sub-Committee of the Committee on Climate Change. Cranfield University, Bedford.
- Grayston SJ, Vaughan D, Jones D (1996) Rhizosphere carbon flow in trees, in comparison with annual plants: the importance of root exudation and its impact on microbial activity and nutrient availability. *Applied Soil Ecology* 5: 29-56.
- Gregorich EG, Rochette P, Hopkins DW, McKim UF, St-Georges P (2006) Tillage-induced environmental conditions in soil and substrate limitation determine biogenic gas production. *Soil Biology and Biochemistry* 38: 2614-2628.
- Grigera MS, Drijber RA, Wienhold BJ (2007) Redistribution of crop residues during row cultivation creates a biologically enhanced environment for soil microorganisms. *Soil Tillage Research* 94:550-554.
- Harding AE, Rivington M, Mineter MJ, Tett SFB (2015) Agro-meteorological indices and climate model uncertainty over the UK. *Climatic Change* 128:113-126.
- Harrison A , Taylor K, Scott A, Poskitt J, Benham D, Grace J, Chaplow J, Rowland P (2008) Potential effects of climate change on DOC release from three different soil types on the Northern Pennines UK: examination using field manipulation experiments. *Global Change Biology* 14:687-702.
- Harrison AM, Plim JFM, Harrison M, Jones LD, Culshaw MG (2012) The relationship between shrink-swell occurrence and climate in southeast England, *Proceedings of the Geologists Association*, vol. 123, no. 4, pp. 556-575.
- Hillier J, Whittaker C, Dailey G et al. (2009) Greenhouse gas emissions from four bioenergy crops in England and Wales: integrating spatial estimates of yield and soil carbon balance in life cycle analyses. *Global Change Biology Bioenergy* 1: 267-281.
- Hobbie SE (1996) Temperature and plant species control over litter decomposition in Alaskan tundra. *Ecological Monographs* 66:503-522.
- Holland JM (2004) The environmental consequences of adopting conservation tillage in Europe: reviewing the evidence. *Agriculture, Ecosystems and Environment* 103: 1-125.
- Holman IP, Nicholls RJ, Berry PM, Harrison PA, Audsley E, Shackley S, Rounsevell MDA (2005) A regional, multi-sectoral and integrated assessment of the impacts of climate and socioeconomic change in the UK. II. Results. *Climatic Change* 71:43-73.
- Hopkins DW, Waite IS, McNicol JW, Poulton PR, Macdonald AJ, O'Donnell AG (2009). Soil organic carbon contents in long-term experimental grassland plots in the UK (Palace Leas and Park Grass) have not changed consistently in recent decades. *Global Change Biology* 15: 1739-1754.

Houghton JT, Callander BA, Varney SK (1992) *Climate Change: The Supplementary Report to the IPCC Scientific Assessment*, Cambridge University Press, Cambridge, UK, 1992.

Hough MN, Jones RJA (1997) The UK Met Office rainfall and evaporation calculation system: MORECS version 2.0 -an overview. *Hydrology and Earth System Sciences* 1:227-239.

Hulme M, Hossell JE, Parry ML (1993). Future climate change and land use in the UK. *Geography Journal* 159:131-147.

Hungate BA, Duval BD, Dijkstra P, Johnson DW, Ketterer ME, Stiling P, Cheng W, Millman J, Hartley A, Stover DB (2014) Nitrogen inputs and losses in response to chronic CO<sub>2</sub> exposure in subtropical oak woodland. *Biogeosciences* 11: 3323-3337.

Ineson P, Benham DG, Poskitt J, Harrison AF, Taylor K, Woods C (1998a) Effects of climate change on nitrogen dynamics in upland soils. 2. A soil warming study. *Global Change Biology* 4:153-61.

Ineson P, Taylor K, Harrison AF, Poskitt J, Benham DG, Tipping E, Woof C (1998b) Effects of climate change on nitrogen dynamics in upland soils. 1. A transplant approach. *Global Change Biology* 4:143-52.

Jackson RB, Sperry JS, Dawson TE (2000) Root water uptake and transport: using physiological processes in global predictions. *Trends in Plant Science* 5: 482-488.

Janssens IA, et al. (2003) Europe's terrestrial biosphere absorbs 7 to 12% of European anthropogenic CO<sub>2</sub> emissions. *Science* 300:1538-1542.

Jastrow JD, Amonette JE, Bailey VL (2007) Mechanisms controlling soil carbon turnover and their potential application for enhancing carbon sequestration. *Climatic Change* 80: 5-23.

Jenny H (1941) *Factors of Soil Formation*. McGraw-Hill, New York, 281 pp.

Jia ZJ, Conrad R (2009) Bacteria rather than Archaea dominate microbial ammonia oxidation in an agricultural soil. *Environmental Microbiology* 11:1658-1671.

Johnson D E, Ward G M, Ramsey JJ (1996) Livestock methane: Current emissions and mitigation potential. In: *Nutrient management of food animals to enhance and protect the environment* (Ed. E. T. Kornegay). Lewis Publishers, New York, NY. 219-234.

IPCC (2013) Summary for Policymakers. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker TF, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Izaurrealde RCC, NJ, Rosenberg RA, Brown JR, Thomson AM (2003) Integrated Assessment of Hadley Centre (HadCM2) Climate-Change Impacts on Agricultural Productivity and Irrigation Water Supply in the Conterminous United States. Part II. Regional Agricultural Production in 2030 and Soil 2095. *Agricultural and Forest Meteorology* 117:97-122.

Murphy JM, Sexton DMH, Jenkins GJ, Booth B, Brown CC, Clark RT, Collins M, Harris GR, Kendon EJ, Betts RA, Brown SJ, Humphrey KA, McCarthy MP, McDonald RE, Stephens A, Wallace C, Warren R, Wilby R, Wood RA (2009) *UK Climate projections science report: climate change projections*. Met Office Hadley Centre, Exeter.

- Kattenberg A, Giorgi F, Grassl H, Meehl GA, Mitchell JFB, Stouffer RJ, Tokioka T, Weaver AJ, and Wigley TML (1996) Climate models - projections of future climate, in *Climate Change 1995: The Science of Climate Change*, edited by J. T. Houghton, L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, pp. 285-357, Cambridge University Press, Cambridge, UK.
- Kay AL, Bell VA, Blyth EM, Crooks SM, Davies HN and Reynard NS (2013) A hydrological perspective on evaporation: Historical trends and future projections in Britain. *Journal of Water and Climate Change* 4: 193-208.
- Kelso BHL, Smith RV, Laughlin RJ, Lennox SD (1997) Dissimilatory nitrate reduction in anaerobic sediments leading to river nitrite accumulation. *Applied and Environmental Microbiology* 63: 4679-4685.
- King JA, Bradley RI and Harrison R (2005) Current trends of soil organic carbon in English arable soils. *Soil Use Management* 21:189-195.
- Kirschbaum MUF (1995) The temperature dependence of soil organic matter decomposition, and the effect of global warming on soil organic C storage. *Soil Biology and Biochemistry* 27: 753-760.
- Kirk GJD, Bellamy PH (2010) Analysis of changes in organic carbon in mineral soils across England and Wales using a simple single-pool model. *European Journal of Soil Science* 61: 406-411.
- Lal R (2008) Carbon sequestration. *Philosophical Transactions of the Royal Society, Series B*, 363: 815-830.
- Leiros MC, Trasar-Cepeda C, Seoane S, Gil-Sotres F (1999) Dependence of mineralization of soil organic matter on temperature and moisture. *Soil Biology Biochemistry* 31: 327-335.
- Liu JX, Zhang DQ, Zhou GY, Faivre-Vuillin B, Deng Q, Wang CL (2008) CO<sub>2</sub> enrichment increases nutrient leaching from model forest ecosystems in subtropical China. *Biogeosciences* 5:1783-1795.
- MacDonald AM, KB Matthews Paterson E, Aspinall RJ (1994) The impact of climate change on the soil-moisture regime of Scottish mineral soils. *Environmental Pollution* 83: 245-250.
- MacDonald NW, Zak DR, Pregitzer KS (1995) Temperature effects on kinetics of microbial respiration and net nitrogen and sulphur mineralization. *Soil Science Society of America Journal* 59: 233-240.
- Machefert SE, Dise NB, Goulding KWT, Whitehead, PG (2002) Nitrous oxide emission from a range of land uses across Europe. *Hydrology and Earth System Sciences*, 6:325-337
- Matzner E, Borken W (2008) Do freeze-thaw events enhance C and N losses from soils of different ecosystems? A review. *European Journal of Soil Science* 59:274-84.
- McCartney AG, Harriman R, Watt AW, Moore DW, Taylor EM, Collen P, Keay EJ (2003) Long-term trends in pH, aluminium and dissolved organic carbon in Scottish fresh waters; implications for brown trout (*Salmo trutta*) survival. *The Science of the Total Environment* 310:133-141.

- Mearns LO (2003) Issues in the impacts of climate variability and change on agriculture. *Climatic Change* 60, 1-6.
- McHugh M (2007) Short-term changes in upland soil erosion in England and Wales: 1999 to 2002. *Geomorphology* 86: 204-213.
- MLCC (2013) Managing the land in a changing climate 2013. Adaptation Sub-Committee Progress Report 2013.
- Moebius BN, van Es HM, Schindelbeck RR, Idowu OJ, Clune DJ, Thies JE (2007) Evaluation of laboratory-measured soil properties as indicators of soil physical quality. *Soil Science* 172:895-912.
- Mohan SB, Schmid M, Jetten M, Cole J (2004) Detection and widespread distribution of the *nrfA* gene encoding nitrite reduction to ammonia, a short circuit in the biological nitrogen cycle that competes with denitrification. *FEMS Microbiology Ecology* 22: 57-63.
- Mojeremane W, Rees RM, Mencuccini M (2010) Effects of site preparation for afforestation on methane fluxes at Harwood Forest, NE England. *Biogeochemistry* 97: 89-107.
- Moore TR, Roulet NT, Waddington JM (1998), Uncertainty in predicting the effect of climatic change on the carbon cycling of Canadian peatlands. *Climatic Change* 40:229- 245.
- Morgan JA, Mosier AR, Milchunas DG, LeCain DR, Nelson JA, Parton W (2004) CO<sub>2</sub> Enhances productivity, alters species composition, and reduces digestibility of shortgrass steppe vegetation. *Ecological Application* 14:208-219.
- Mullan D, Favis-Mortlock D, Fealy R (2012) Addressing key limitations associated with modelling soil erosion under the impacts of future climate change. *Agricultural and Forest Meteorology* 156:18-30.
- Murphy JM, Sexton DMH, Jenkins GJ, Booth B, Brown CC, Clark RT, Collins M, Harris GR, Kendon EJ, Betts RA, Brown SJ, Humphrey KA, McCarthy MP, McDonald RE, Stephens A, Wallace C, Warren R, Wilby R, Wood RA (2009) UK Climate projections science report: climate change projections. Met Office Hadley Centre, Exeter.
- Naden PS, Watts CD (2001) Estimating climate-induced change in soil moisture at the landscape scale: an application of five areas of ecological interest in the UK. *Climatic Change* 49: 411-440.
- Nearing MA, Jetten V, Cerdan O, Couturier A, Hernandez M, Le Bissonnais Y, Nicholas MH, Nunes JP, Renschler CS, Souchere V, Van Oost K (2005) Modeling response of soil erosion and runoff to changes in precipitation and cover. *Catena* 61: 131-154.
- Nielsen U, Ayres E, Wall D, Bardgett RD (2011) Soil biodiversity and carbon cycling: a review and synthesis of studies examining diversity-function relationships. *European Journal of Soil Science* 62: 105-116.
- Olesen JE, Carter TR, Diaz-Ambrona CH, Fronzek S, Heidmann T, Hickler T, et al. (2007) Uncertainties in projected impacts of climate change on European agriculture and terrestrial ecosystems based on scenarios from regional climate models. *Climatic Change* 81:123-43.

- Osborne B, Saunders M, Walmsley D, Jones M, Smith P (2010) Key questions and uncertainties associated with the assessment of the cropland greenhouse gas balance, *Agriculture, Ecosystems and Environment* 139: 293-301.
- Passioura JB (1991). Soil structure and plant growth. *Australian Journal of Soil Research* 29: 717-728
- Pawson RR, Lord DR, Evans MG, Allott TEH (2008) Fluvial organic carbon flux from an eroding peatland catchment, southern Pennines, UK. *Hydrology and Earth System Sciences*, 12: 625-634
- Phillips RP, Finzi AC, Bernhardt ES (2011) Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO<sub>2</sub> fumigation. *Ecology Letters* 14:187-194.
- Phillips RP, Meier IC, Bernhardt ES, Grandy AS, Wickings K, Finzi AC (2012) Roots and fungi accelerate carbon and nitrogen cycling in forests exposed to elevated CO<sub>2</sub>. *Ecology Letters* 15:1042-1049.
- Pierce FJ, Nowak P (1999) Aspects of precision agriculture. *Advances in Agronomy* 67:1-85.
- Pignatelli M, De Nobili M (1993) Relationship between soil porosity and root development and soil enzyme activity in cultivated soils. *Geoderma* 56:243–256
- Porporato A, Daly E, Rodriguez-Iturbe I (2005) Soil water balance and ecosystem response to climate change. *Am Nat* 164:625-632
- Post WM, Kwon KC (2000) Soil carbon sequestration and land-use change: processes and potential. *Global Change Biology* 6: 317-327.
- Powlson D, Stirling C, Jat M, Gerard B, Palm C, Sanchez P, Cassman K (2014) Limited potential of no-till agriculture for climate change mitigation. *Nature Climate Change* 4, 678-683.
- Prescott CE (2009) Litter decomposition: what controls it and how can we alter it to sequester more carbon in forest soils? *Biogeochemistry* 101:133-149.
- Rees RM, Bingham IJ, Baddeley JA, Watson CA (2005) The role of plants and land management in sequestering soil carbon in temperate arable and grassland ecosystems. *Geoderma* 128:130-154.
- Qafoku NP (2014) Overview of different aspects of climate change effects on soils. Prepared for the U.S. Department of Energy under Contract DE-AC05-76RL01830. <http://www.Ntis.gov>.
- Reeve and Fausey (1974) Drainage timeliness of farming operations. *Drainage for agriculture. Monograph 17 (ed). Madison Wisconsin: American Society of Agronomy* pp 55-66.
- Reicosky DC, Archer DW (2007) Moldboard plow tillage depth and short-term carbon dioxide release. *Soil Tillage Research* 94:109-121.
- Reinds GJ, Posch M, Leemans R (2009) Modelling recovery from soil acidification in



European forests under climate change. *Science of the Total Environment* 407: 5663-5673.

Reynolds WD, Drury CF, Tan CS, Fox CA, Yang XM (2009) Use of indicators and pore volume function characteristics to quantify soil physical quality. *Geoderma* 152:252-263.

Reynolds B, Chamberlain PM, Poskitt J et al. (2013) Countryside survey: national 'soil change' 1978–2007 for topsoils in Great Britain - acidity, carbon, and total nitrogen status. *Vadose Zone Journal*, 12, Available at: <https://www.soils.org/publications/citation-manager/vzj/12/2/vzj2012.0114>.

Rimal BK, Lal R (2009) Soil and carbon losses from five different land management areas under simulated rainfall. *Soil Tillage Research* 106:62-70.

Rivett MO, Smith JNW, Buss SR, Morgan P (2007) Nitrate occurrence and attenuation in the major aquifers of 523 England and Wales. *Quarterly Journal of Engineering Geology and Hydrogeology* 40: 335-352.

Rounsevell MDA, Brignall AP and Siddons PA (1996). Potential climate change effects on the distribution of agricultural grassland in England and Wales. *Soil Use and Management* 12: 31-44.

Rounsevell M, Evans SP, Bullock P (1999) Climate change and agricultural soils: impacts and adaptation. *Climatic Change* 43:683-709.

Rosenzweig C, Hillel D (1998) *Climate Change and the Global Harvest. Potential Impacts of the Greenhouse Effect on Agriculture*. Oxford University Press, Inc., New York.

Rowe EC, Emmett BA, Frogbrook ZL, Robinson DA, Hughes S (2012) Nitrogen deposition and climate effects on soil nitrogen availability: Influences of habitat type and soil characteristics. *Science of the Total Environment* 434: 62-70.

Rustad LE, Campbell JL, Marion GM, Norby RJ, Mitchell MJ, Hartley AE, Cornelissen JHC, Gurevitch J (2001) A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia* 126:543-562.

Schils R, Kuikman P, Liski J, Van Oijen M, Smith P, Webb J, Hiederer R (2008) Review of existing information on the interrelations between soil and climate change. European Communities, 2008 (ClimSoil). Final report. Technical report, doi:10.2779/12723.

Schimel DS (1995) Terrestrial biogeochemical cycles: global estimates with remote sensing. *Remote Sensing of Environment* 51: 49-56.

Schlesinger WH (1995) Soil respiration and changes in soil carbon stocks. Pages 159-168 in: Woodwell GM, Mackenzie FT. eds. *Biotic Feedbacks in the Global Climate System: Will the Warming Feed the Warming?* New York, Oxford University Press.

Schmidt IK, Emmett BA, Tietema A et al (2004) Soil solution chemistry and element fluxes in three European heathlands and their responses to warming and drought. *Ecosystems* 7: 638- 649.

Schulze ED, Freibauer A (2005) Carbon unlocked from soils. *Nature* 437: 245-248.

Sgouridis F, Ullah S (2014) Denitrification potential of organic, forest and grassland soils in

- the Ribble-Wyre and Conwy River catchments, UK. *Environmental Science: Processes Impacts* 16: 1551-1562.
- Sheffield J, Wood EF and Roderick ML (2012) Little change in global drought over the past 60 years. *Nature* 491: 435-438.
- Shepherd M, Pearce B, Cormack B, Phillipps L, Cuttle S, Bhogal A, Costagin P, Unwin R (2002) An assessment of the environmental impacts of organic farming. A review for Defra-funded project OF0405 - Defra.
- Shiels AB (2006) Leaf litter decomposition and substrate chemistry of early successional species on landslides in Puerto Rico. *Biotropica* 38:348-53.
- Six J, Bossuyt H, Degryze S, Denef K (2004) A history of research on the link between (micro) aggregates, soil biota, and soil organic matter dynamics. *Soil and Tillage Research* 79: 7-31.
- Skiba U, Jones SK, Dragosits U, Drewer J, Fowler D, Rees RM, Pappa VA, Cardenas L, Chadwick D, Yamulki S, Manning AJ (2012) UK emissions of the greenhouse gas nitrous oxide. *Philosophical Transactions of the Royal Society B* 367:1175-1185.
- Skj  th CA, Geels C (2013) The effect of climate and climate change on ammonia emissions in Europe, *Atmos. Chemical Physics* 13:117-128.
- Smith KA, Ball T, Conen F, Dobbie KE, Massheder J, Rey A (2003) Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes. *European Journal of Soil Science* 54:779-791.
- Smith P, Smith J, Wattenbach M, Meyer J, Lindner M, Zaehle S, Hiederer R, Jones RJA, Montanarella L, Rounsevell M, Reginster I, Kankaanp    S (2006) Projected changes in mineral soil carbon of European forests, 1990-2100. *Canadian Journal of Soil Science* 86: 159-169.
- Smith P, Chapman SJ, Scott WA, Black HJ, Wattenbach M, Milne R, Campbell CD, Lilly, A, Ostle N, Levy PE, Lumsdon DG, Millard P, Towers W, Zaehle S, Smith JU (2007) Climate change cannot be entirely responsible for soil carbon loss observed in England and Wales, 1978-2003. *Global Change Biology* 13: 2605-2609.
- Smith P (2012) Soils and climate change. *Current Opinion in Environmental Sustainability* 4:539-544.
- Smith SE, Read DJ (1997) *Mycorrhizal Symbiosis*. Academic Press, New York, pp. 379-409.
- Smith SV, Renwick WH, Buddemeier RW, Crossland CJ (2001) Budgets of soil erosion and deposition for sediments and sedimentary organic carbon across the conterminous United States. *Global Biogeochemical Cycles* 15: 697-707.
- Stanford G, Smith SJ (1972) Nitrogen mineralization potentials of soils. *Soil Science Society of America Proceedings* 36:465-72.
- Stopes C, Lord EI, Philipps L, Woodward L (2002) Nitrate leaching from organic farms and conventional farms following best practice. *Soil Use and Management* 18: 256-264.
- Stopnisek N., Gubry-Rangin C., H        S., Nicol G. W., Mandic-Mulec I., Prosser J. I. (2010). Thaumarchaeal ammonia oxidation in an acidic forest peat soil is not influenced by ammonium amendment. *Applied Environmental Microbiology* 76: 7626-7634.

- Stuart ME, Chilton PJ, Kinniburgh DG, Cooper DM (2007) Screening for long-term trends in groundwater nitrate monitoring data. *Quarterly Journal of Engineering Geology and Hydrogeology* 40:361-76.
- Stuart ME, Gooddy DC, Bloomfield JP, Williams AT (2011) Future impact may also be driven by economic responses to climate change. *Science of the Total Environment* 409: 2859-2873.
- Tate RL (1992) *Soil Organic Matter: Biological and Ecological Effects* Krieger. Publishing Company, Florida, USA.
- Thomson AM (2008) Inventory and projections of the UK emissions by sources and removal by sinks due to land use, land use change and forestry (Ed.). London. Defra climate. Energy and ozone. Science and analysis division.
- Thomsen IK (2005) Crop N utilization and leaching losses as affected by time and method of application of farmyard manure. *European Journal of Agronomy* 22: 1-9.
- Van Veen JA, Liljeroth E, Lekkerkerk LJA, Van De Geijn SC (1991) Carbon fluxes in plant-soil systems at elevated atmospheric carbon dioxide levels. *Ecological Applications* 1:175-181.
- Vinther FP, Dahlmann-Hansen L (2005) Effects of ridging on crop performance and symbiotic N<sub>2</sub> fixation in fababean (*Vicia faba* L.). *Soil Use and Management* 21:205-211.
- Waddington JM, Day SM (2007) Methane emissions from a peatland following restoration. *Journal of Geophysical Research: Biogeosciences* 112: G03018.
- Webb J, Loveland PJ, Chambers BJ, Mitchell R Garwood T (2001). The impact of modern farming practices on soil fertility and quality in England and Wales. *Journal of Agricultural Science* 137:127-138.
- West TO, Post WM (2002) Soil organic carbon sequestration rates by tillage and crop rotation: a global data analysis. *Soil Science Society of America Journal* 66: 1930-1946.
- White RE, Wellings SR, Bell JP (1983) Seasonal variations in nitrate leaching in structured clay soils under mixed land use. *Agricultural Water Management* 7: 391-410.
- Williams JR, Nearing MA, Nicks A, Skidmore E, Valentine C, King K, Savabi R (1996) Using soil erosion models for global change studies. *Journal of Soil and Water Conservation* 51: 381-385.
- Worrall F, Burt T, Shedden R (2003) Long term records of riverine dissolved organic matter. *Biogeochemistry* 64:165-178.
- Worrall F, Burt T, Adamson JK (2004) Can climate change explains increases in DOC flux from upland catchments? *Science of the Total Environment* 326: 95-112.
- Worrall F, Burt TP (2004) Time series analysis of long term river DOC records. *Hydrological Processes* 18:893-911.

- Yin HJ, Li YF, Xiao J, Xu ZF, Cheng XY, Liu Q (2013) Enhanced root exudation stimulates soil nitrogen transformations in a subalpine coniferous forest under experimental warming. *Global Change Biology* 19:2158-2167.
- Zak DR, Pregitzer KS, King JS, Holmes WE (2000) Elevated atmospheric CO<sub>2</sub>, fine roots and the response of soil microorganisms: a review and hypothesis. *New Phytologist* 147:201-222.
- Zhalnina K, de Quadros PD, Camargo FAO, Triplett EW (2012) Drivers of archaeal ammonia-oxidizing communities in soil. *Frontiers in Microbiology* 3:210.
- Zhang D, Hui D, Luo Y, Zhou G (2008) Rates of litter decomposition in terrestrial ecosystems: global patterns and controlling factors. *Journal of Plant Ecology* 1:1-9.